

Package ‘ljr’

May 1, 2016

Version 1.4-0

Date 2016-4-30

Title Logistic Joinpoint Regression

Author Michal Czajkowski, Ryan Gill, Greg Rempala

Maintainer Ryan Gill <ryan.gill@louisville.edu>

Description Fits and tests logistic joinpoint models.

License GPL (>= 2)

NeedsCompilation yes

Repository CRAN

Date/Publication 2016-05-01 18:55:59

R topics documented:

kcm	2
ljr0	2
ljr01	3
ljr1	5
ljr11	6
ljrb	7
ljrf	8
ljrk	9
ljrk	10
ljrkk	12
Index	14

kcm

*Kentucky yearly cancer mortality from 1999-2005.***Description**

This table gives the yearly mortality counts due to neoplasms (ICD 10 codes C00-D48) and population sizes for Kentucky from 1999-2005. For more information, see <http://wonder.cdc.gov/wonder/help/cmfm.html>.

Usage

```
data(kcm)
```

Format

A 7 by 3 data frame.

Source

Centers for Disease Control and Prevention, National Center for Health Statistics. Compressed Mortality File 1999-2005. CDC WONDER On-line Database, compiled from Compressed Mortality File 1999-2005 Series 20 No. 2K, 2008. Accessed at <http://wonder.cdc.gov/cmfm-icd10.html> on May 5, 2008.

ljr0

*MLE with 0 joinpoints***Description**

Determines the maximum likelihood estimate of model coefficients in the logistic joinpoint regression model with no joinpoints.

Usage

```
ljr0(y,n,tm,X,ofst)
```

Arguments

y	the vector of Binomial responses.
n	the vector of sizes for the Binomial random variables.
tm	the vector of observation times.
X	a design matrix containing other covariates.
ofst	a vector of known offsets for the logit of the response.

Details

The re-weighted log-likelihood is the log-likelihood divided by the largest component of n .

Value

Coef	A table of coefficient estimates.
wlik	The maximum value of the re-weighted log-likelihood.

Author(s)

The authors are Michal Czajkowski, Ryan Gill, and Greg Rempala. The software is maintained by Ryan Gill <rsgill01@louisville.edu>.

References

Czajkowski, M., Gill, R. and Rempala, G. (2008). Model selection in logistic joinpoint regression with applications to analyzing cohort mortality patterns. *Statistics in Medicine* 27, 1508-1526.

See Also

[ljr01](#), [ljrb](#), [ljrf](#)

Examples

```
data(kcm)
attach(kcm)
ljr0(Count,Population,Year+.5)
```

ljr01	<i>Perform test of 0 vs 1 joinpoints.</i>
-------	---

Description

This function tests the null hypothesis of 0 joinpoints versus the alternative of one joinpoint based on the likelihood ratio test statistic. The p-value is determined by a Monte Carlo method.

Usage

```
ljr01(y, n, tm, X, ofst, R=1000, alpha=.05)
```

Arguments

y	the vector of Binomial responses.
n	the vector of sizes for the Binomial random variables.
tm	the vector of ordered observation times.
X	a design matrix containing other covariates.
ofst	a vector of known offsets for the logit of the response.
R	number of Monte Carlo simulations.
alpha	significance level of the test.

Details

The re-weighted log-likelihood is the log-likelihood divided by the largest component of n.

Value

pval	The estimate of the p-value via simulation.
Coef	A table of coefficient estimates.
Joinpoint	The estimates of the joinpoint, if it is significant.
wlik	The maximum value of the re-weighted log-likelihood.

Author(s)

The authors are Michal Czajkowski, Ryan Gill, and Greg Rempala. The software is maintained by Ryan Gill <rsgill01@louisville.edu>.

References

Czajkowski, M., Gill, R. and Rempala, G. (2008). Model selection in logistic joinpoint regression with applications to analyzing cohort mortality patterns. *Statistics in Medicine* 27, 1508-1526.

See Also

[ljr0](#), [ljr1](#)

Examples

```
data(kcm)
attach(kcm)
set.seed(12345)
## Not run: ljr01(Count,Population,Year+.5,R=20)
```

ljr1	<i>MLE with 1 joinpoint</i>
------	-----------------------------

Description

Determines the maximum likelihood estimates of model coefficients in the logistic joinpoint regression model with one joinpoint.

Usage

```
ljr1(y, n, tm, X, ofst, summ=TRUE)
```

Arguments

y	the vector of Binomial responses.
n	the vector of sizes for the Binomial random variables.
tm	the vector of ordered observation times.
X	a design matrix containing other covariates.
ofst	a vector of known offsets for the logit of the response.
summ	a boolean indicator of whether summary tables should be returned.

Details

The re-weighted log-likelihood is the log-likelihood divided by the largest component of n.

Value

Coef	A table of coefficient estimates.
Joinpoint	The estimate of the joinpoint.
wlik	The maximum value of the re-weighted log-likelihood.

Author(s)

The authors are Michal Czajkowski, Ryan Gill, and Greg Rempala. The software is maintained by Ryan Gill <rsgill01@louisville.edu>.

References

Czajkowski, M., Gill, R. and Rempala, G. (2008). Model selection in logistic joinpoint regression with applications to analyzing cohort mortality patterns. *Statistics in Medicine* 27, 1508-1526.

See Also

[ljr01](#), [ljrb](#), [ljrf](#)

Examples

```
data(kcm)
attach(kcm)
ljr1(Count,Population,Year+.5)
```

ljr11

Test coefficients conditioned on K=1 joinpoint.

Description

This function performs the likelihood ratio tests to find p-values in testing the significance of each of the coefficients as well as the intercept and ordered observation times. The p-values are determined by a Monte Carlo method.

Usage

```
ljr11(y,n,tm,X,ofst,R=1000)
```

Arguments

y	the vector of Binomial responses.
n	the vector of sizes for the Binomial random variables.
tm	the vector of ordered observation times.
X	a design matrix containing other covariates.
ofst	a vector of known offsets for the logit of the response.
R	number of Monte Carlo simulations.

Details

The re-weighted log-likelihood is the log-likelihood divided by the largest component of n.

Value

pvals	The estimates of the p-values via simulation.
-------	---

Author(s)

The authors are Michal Czajkowski, Ryan Gill, and Greg Rempala. The software is maintained by Ryan Gill <rsgill01@louisville.edu>.

References

Czajkowski, M., Gill, R. and Rempala, G. (2008). Model selection in logistic joinpoint regression with applications to analyzing cohort mortality patterns. *Statistics in Medicine* 27, 1508-1526.

See Also[ljr1](#)**Examples**

```

data(kcm)
attach(kcm)
set.seed(12345)
## Not run: ljr11(Count,Population,Year+.5,R=20)

```

ljrb	<i>Perform backward joinpoint selection algorithm with upper bound K.</i>
------	---

Description

This function performs the backward joinpoint selection algorithm with K maximum possible number of joinpoints based on the likelihood ratio test statistic. The p-value is determined by a Monte Carlo method.

Usage

```
ljrb(K, y, n, tm, X, ofst, R=1000, alpha=.05)
```

Arguments

K	the pre-specified maximum possible number of joinpoints
y	the vector of Binomial responses.
n	the vector of sizes for the Binomial random variables.
tm	the vector of ordered observation times.
X	a design matrix containing other covariates.
ofst	a vector of known offsets for the logit of the response.
R	number of Monte Carlo simulations.
alpha	significance level of the test.

Details

The re-weighted log-likelihood is the log-likelihood divided by the largest component of n.

Value

pvals	The estimates of the p-values via simulation.
Coef	A table of coefficient estimates.
Joinpoints	The estimates of the joinpoint, if it is significant.
wlik	The maximum value of the re-weighted log-likelihood.

Author(s)

The authors are Michal Czajkowski, Ryan Gill, and Greg Rempala. The software is maintained by Ryan Gill <rsgill01@louisville.edu>.

References

Czajkowski, M., Gill, R. and Rempala, G. (2008). Model selection in logistic joinpoint regression with applications to analyzing cohort mortality patterns. *Statistics in Medicine* 27, 1508-1526.

See Also

[ljrk](#), [ljrf](#)

Examples

```
data(kcm)
attach(kcm)
set.seed(12345)
## Not run: ljrb(1,Count,Population,Year+.5,R=20)
```

ljrf	<i>Perform forward joinpoint selection algorithm with unlimited upper bound.</i>
------	--

Description

This function performs the full forward joinpoint selection algorithm based on the likelihood ratio test statistic. The p-value is determined by a Monte Carlo method.

Usage

```
ljrf(y,n,tm,X,ofst,R=1000,alpha=.05)
```

Arguments

y	the vector of Binomial responses.
n	the vector of sizes for the Binomial random variables.
tm	the vector of ordered observation times.
X	a design matrix containing other covariates.
ofst	a vector of known offsets for the logit of the response.
R	number of Monte Carlo simulations.
alpha	significance level of the test.

Details

The re-weighted log-likelihood is the log-likelihood divided by the largest component of n.

Value

pvals	The estimates of the p-values via simulation.
Coef	A table of coefficient estimates.
Joinpoints	The estimates of the joinpoint, if it is significant.
wlik	The maximum value of the re-weighted log-likelihood.

Author(s)

The authors are Michal Czajkowski, Ryan Gill, and Greg Rempala. The software is maintained by Ryan Gill <rsgill01@louisville.edu>.

References

Czajkowski, M., Gill, R. and Rempala, G. (2008). Model selection in logistic joinpoint regression with applications to analyzing cohort mortality patterns. *Statistics in Medicine* 27, 1508-1526.

See Also

[ljrk](#), [ljrb](#)

Examples

```
data(kcm)
attach(kcm)
set.seed(12345)
## Not run: ljrjk(Count,Population,Year+.5,R=20)
```

ljrjk	<i>Perform test of j vs k joinpoints.</i>
-------	---

Description

This function tests the null hypothesis of j joinpoint(s) versus the alternative of k joinpoint(s) based on the likelihood ratio test statistic. The p-value is determined by a Monte Carlo method.

Usage

```
ljrjk(j,k,y,n,tm,X,ofst,R=1000,alpha=.05)
```

Arguments

j,k	pre-specified number of joinpoints in the null and alternative hypotheses (the smaller is used for the null).
y	the vector of Binomial responses.
n	the vector of sizes for the Binomial random variables.
tm	the vector of ordered observation times.

X	a design matrix containing other covariates.
ofst	a vector of known offsets for the logit of the response.
R	number of Monte Carlo simulations.
alpha	significance level of the test.

Details

The re-weighted log-likelihood is the log-likelihood divided by the largest component of n.

Value

pval	The estimate of the p-value via simulation.
Coef	A table of coefficient estimates.
Joinpoint	The estimates of the joinpoint, if it is significant.
wlik	The maximum value of the re-weighted log-likelihood.

Author(s)

The authors are Michal Czajkowski, Ryan Gill, and Greg Rempala. The software is maintained by Ryan Gill <rsgill01@louisville.edu>.

References

Czajkowski, M., Gill, R. and Rempala, G. (2008). Model selection in logistic joinpoint regression with applications to analyzing cohort mortality patterns. *Statistics in Medicine* 27, 1508-1526.

See Also

[ljrk](#)

Examples

```
data(kcm)
attach(kcm)
set.seed(12345)
## Not run: ljrk(0,1,Count,Population,Year+.5,R=20)
```

ljrk

MLE with k joinpoints

Description

Determines the maximum likelihood estimates of model coefficients in the logistic joinpoint regression model with k joinpoints.

Usage

```
ljrk(k, y, n, tm, X, ofst)
```

Arguments

k	the pre-specified number of joinpoints (with unknown locations).
y	the vector of Binomial responses.
n	the vector of sizes for the Binomial random variables.
tm	the vector of ordered observation times.
X	a design matrix containing other covariates.
ofst	a vector of known offsets for the logit of the response.

Details

The re-weighted log-likelihood is the log-likelihood divided by the largest component of n.

Value

Coef	A table of coefficient estimates.
Joinpoints	The estimates of the joinpoints.
wlik	The maximum value of the re-weighted log-likelihood.

Author(s)

The authors are Michal Czajkowski, Ryan Gill, and Greg Rempala. The software is maintained by Ryan Gill <rsgill01@louisville.edu>.

References

Czajkowski, M., Gill, R. and Rempala, G. (2008). Model selection in logistic joinpoint regression with applications to analyzing cohort mortality patterns. *Statistics in Medicine* 27, 1508-1526.

See Also

[ljrb](#), [ljrf](#)

Examples

```
data(kcm)
attach(kcm)
ljrk(1, Count, Population, Year+.5)
```

`ljrkk`*Test coefficients conditioned on $K=k$ joinpoint.*

Description

This function performs the likelihood ratio tests to find p-values in testing the significance of each of the coefficients as well as the intercept and ordered observation times. The p-values are determined by a Monte Carlo method.

Usage

```
ljrkk(k, y, n, tm, X, ofst, R=1000)
```

Arguments

<code>k</code>	the pre-specified number of joinpoints (with unknown locations).
<code>y</code>	the vector of Binomial responses.
<code>n</code>	the vector of sizes for the Binomial random variables.
<code>tm</code>	the vector of ordered observation times.
<code>X</code>	a design matrix containing other covariates.
<code>ofst</code>	a vector of known offsets for the logit of the response.
<code>R</code>	number of Monte Carlo simulations.

Details

The re-weighted log-likelihood is the log-likelihood divided by the largest component of `n`.

Value

`pvals` The estimates of the p-values via simulation.

Author(s)

The authors are Michal Czajkowski, Ryan Gill, and Greg Rempala. The software is maintained by Ryan Gill <rsgill01@louisville.edu>.

References

Czajkowski, M., Gill, R. and Rempala, G. (2008). Model selection in logistic joinpoint regression with applications to analyzing cohort mortality patterns. *Statistics in Medicine* 27, 1508-1526.

See Also

[ljrk](#)

Examples

```
data(kcm)
attach(kcm)
set.seed(12345)
## Not run: ljrkk(1,Count,Population,Year+.5,R=20)
```

Index

*Topic **datasets**

kcm, [2](#)

*Topic **nonlinear**

ljr0, [2](#)

ljr01, [3](#)

ljr1, [5](#)

ljr11, [6](#)

ljrb, [7](#)

ljrf, [8](#)

ljrjk, [9](#)

ljrk, [10](#)

ljrkk, [12](#)

kcm, [2](#)

ljr0, [2, 4](#)

ljr01, [3, 3, 5](#)

ljr1, [4, 5, 7](#)

ljr11, [6](#)

ljrb, [3, 5, 7, 9, 11](#)

ljrf, [3, 5, 8, 8, 11](#)

ljrjk, [9](#)

ljrk, [8-10, 10, 12](#)

ljrkk, [12](#)